

Investigation of Seed Germination and Seedling Characteristics of Safflower Variety Under Salt Stress Conditions

Hossein Sadeghi Give¹, Morteza Firoozi², Negin Ahmadzadeh³

1, 2 Department of Agriculture, Birjand branch, Islamic Azad University, Birjand, Iran.

3. Department of Agronomy and Plant Breeding, College of Agriculture, Karaj branch, Islamic Azad University, Alborz, Iran.

Corresponding author email: hsadeghi@iaubir.ir

ABSTRACT: In order to study the effect of salinity stress on germination and early seedling growth of six safflower genotypes namely FERAMAN, SINA, KM12, KM19, SIRIN and Kose by using five concentrations of NaCl (0, -0.3, -0.5, -1 and -1.5 MPa) a factorial experiment was designed using Completely Randomized Design (CRD) with three replications in. Results of ANOVA showed that salt stress adversely affected the germination percentage, germination rate; shoot length, root length, seedling length, root/shoot length ratio, seed vigour, and germination index and mean germination time of all 6 genotypes of safflower, which demonstrates high diversity among genotypes that enabled us to screen salinity tolerant cultivar. At the highest salt level (-1.5 MPa), Kose produced maximum germination percentage and germination rate of all genotypes and they were considered as relatively tolerant. Best level of NaCl concentration in root length, shoot length, seedling length and seed vigour was -0.3 MPa. Seed vigour increased with increase in osmotic potential until -0.3 MPa but decreased in -0.5 MPa. Results of cluster analysis (Ward's minimum variance method) at the highest salt level (-1.5 MPa) classified all genotypes into three group. According to the obtained results, we found that Kose is the most resistant and FERAMAN, SINA and SIRIN are the most sensitive genotypes.

Keywords: Cluster analysis, Germination indices, NaCl, Seed vigour

INTRODUCTION

Among various environmental stresses, soil salinity has become a critical problem worldwide due to its dramatic effects on plant physiology and performance (Golbashy *et al.*, 2010). Salinity in soil or water is one of the major stresses and especially in arid and semi arid regions, can severely limit crop production (Shannon, 1998). Breeders seek to develop and identify cultivars that are more tolerant of salinity and water stress (Janmohammadi *et al.*, 2008). Germination is generally considered to be the developmental stage that is most salt-sensitive, especially for crops exposed to hostile environments (Ashraf and Wahid, 2000).

Salinity impairs seed germination, reduces nodule formation, retards plant development and reduces crop yield (Greenway and Munns, 1980). Soil salinity may affect the germination of seeds either by creating osmotic potential external to the seeds preventing water uptake or through the toxic effects of Na⁺ and Cl⁻ ions on germinating seed (Golbashy *et al.*, 2010; Khajeh-Hosseini *et al.*, 2003; Atak *et al.*, 2006; Kaya *et al.*, 2006). Salinity delays the onset, reduces the rate and increases the dispersion of germination events, resulting in reduced plant growth and final crop yield (Ashraf and Foolad, 2005). Absence of germination in salinity soil is very often due to the high concentration of salt in the soil where the seeds are sown. The reason is that the salt solution moves upward, following the evaporation at soil level (Bernstein, 1974). Salt disturbs both germination and plant growth (Fowler, 1991). The main salt-induced physiological disorder is diminished seed imbibitions because of the low solute potential within the saline growth medium (Debez *et al.*, 2004). Seed may be more sensitive to stress than mature plants because of exposure the dynamic environment close to the soil surface. One of the commonest experiments in germination of the seeds is the application of NaCl. Seed response to salinity can be simulated by NaCl induced ionic stress in the germination experiments. Ionic stress is caused by a toxic accumulation of NaCl in plant tissues. Germination rates decrease with an increase in NaCl concentration (Murillo-Amador, *et al.* 2002).

Thus, the salt-affected soils can be utilized by growing salt tolerant plants, whether halophytes or crops (Siddiqi *et al.* 2007). With this fact in mind, it is imperative to explore intra-specific (inter-cultivar) variation for

salt tolerance of a crop by screening its available germplasm. For instance, a great magnitude of inter-cultivar variation for salt tolerance has been observed in different species such as wheat (Ashraf & McNeilly, 1988), lentil (Ashraf & Waheed, 1993), barley (Belkhdja *et al.*, 1994), cotton (Ashraf & Ahmad, 1999), Brassica napus (Ulfat *et al.*, 2007) and Safflower (Siddiqi *et al.* 2007). Safflower (*Carthamus tinctorius* L.) is one of the prospective oil-seed crops, because it yields about 32-40% seed oil (Weiss, 1983). However, due to its considerable salt tolerance compared with commonly grown oil-seed crops, it is usually cultivated in arid and semi-arid regions where soil salinity is one of the major threats to agriculture (Kaya, 2009).

The research has shown that in response to soil salinity, seedlings growth, leaves area, root biomass and shoot biomass have all been reduced (Redmann *et al.*, 1994). Although salt stress adversely affects the growth of safflower plants at all developmental stages (Kaya *et al.*, 2003; Jamil *et al.*, 2006; Golbashy *et al.*, 2010), varietal differences in salt tolerance of safflower have been observed at germination (Ghorashy *et al.*, 1972), at adult (Ashraf & Fatima, 1995) as well as at both germination and adult growth stages (Francois & Bernstein, 1964). However, Kaya *et al.* (2006) reported that germination percentage was not influenced by NaCl level of 23.5 $ds\ m^{-1}$. Mohammed *et al.* (2002) reported that by NaCl levels germination percentage decreased and mean germination time increased proportionately.

The present study was therefore, conducted with the objectives to determine the response of safflower genotype to salinity stress at germination and seedling stages under controlled conditions. Moreover, NaCl was used for salinity stress induction in safflower.

MATERIALS AND METHODS

In order to study the effects of salinity stress on germination and early seedling growth in safflower genotypes, an experiment was conducted in factorial form, using a completely randomized design with three replications. In this experiment, six safflower genotypes inclusive FERAMAN, SINA, KM12, KM19, SIRIN and Kose were evaluated in five levels of salinity treatment (distilled water as control, -0.3, -0.5, -1 and -1.5 MPa) by using different NaCl concentrations. This experiment was carried out at Agronomy laboratory, Islamic Azad University- Birjand Branch.

The seeds were sterilized by soaking in a 5% solution of hypochlorite sodium for 5 min. After the treatment, the seeds were washed several times with distilled water. 25 seeds were put in each Petridish (with 9cm diameter) on filter paper moistened with respective treatment in 3 replications. The petridishes were covered to prevent the loss of moisture by evaporation. The petridishes were put into an incubator for 12 days at 25 centigrade degrees temperature and 65% relative humidity. Every 24 hours after soaking, germination percentage and other traits were recorded daily. After 12 days of incubation, shoot length, root length, seed vigour and root to shoot ratio of germinated seeds was measured. Seeds were considered germinated when the emergent radical reached 2 mm length. Germination percentage, germination Rate and seed vigour were calculated using the following formulas:

$$\text{Formula 1: } GP = (SNG/SNO) \times 100 \%$$

Where GP is germination percentage, SNG is the number of germinated seeds, and SNO is the number of experimental seeds with viability (Close and Wilson 2002; Danthu *et al.* 2003).

$$\text{Formula 2: } GR = \sum N / \sum (n \times g)$$

Where: GR: Germination rate; n: number of germinated seed on gth day and g: Number of total germinated seeds

$$\text{Formula 3: Seed Vigour} = [\text{seedling length (cm)} \times \text{germination percentage}]$$

Analysis of variance was performed using standard techniques and differences between the means were compared through Duncan multiple range test ($P < 0.05$) using SAS release 9.1 (SAS, 2002) software package. All investigated traits were subjected to hierarchical cluster analysis using procedure Ward's minimum variance method as a clustering algorithm using Stat Graphics Plus (Ver 2.1) software. Ward's minimum method is a hierarchical clustering procedure in which similarity used to join clusters is calculated as the sum of squares between the two clusters summed over all variables (Hair *et al.*, 1998). It minimizes them within cluster sums of squares across all partitions.

RESULT

Analysis of variance showed that, there were significantly difference between genotypes, salinity stress levels and them interaction. The results of this study reveal that various concentrations of NaCl had a significant effect on the all measured traits (Table 1). The control showed clear genetically differences among the genotypes regards germination percentage, and such differences were statistically significant.

Germination percentage of all safflower genotypes was adversely affected due to the application of different levels (0, -0.3, -0.5, -1 and -1.5 MPa) of NaCl.

Table 1. Analysis of variance of measured traits of safflower genotypes under salinity stress

S.O.V	df	Germination Percentage	Germination Rate	Root Length (mm)	Shoot Length (mm)	Seedling Length (cm)
Genotype	5	2425.23**	1987.12**	14.09**	29.61**	68.35**
stress	4	690.74**	1139.94**	28.32**	58.19**	186.83**
Genotypexstress	20	179.54*	88.33**	3.87**	4.14**	11.88**
error	60	82.94	19.82	1.13	0.83	2.03

Table 1 . continued-

S.O.V	df	Seed Vigour	Root/Shoot Length (mm)	Germination Index	Mean Germination Time (day)
Genotype	5	711732.29**	2.12**	3594.79**	2.59**
stress	4	1278344.18**	0.03ns	1143.17**	1.45**
Genotypexstress	20	94812.71**	0.15ns	120.12**	0.04ns
error	60	21757.84	0.13	43.72	0.02

*, **, ns: significant at 5%, 1% level and not significant, respectively

Also analysis of variance showed that, interaction effects was significant for all investigated characters except root to shoot length ratio and mean germination time.

The differences between the means (Genotypes and salinity stress levels) were compared by Duncan multiple range test and are shown in Table 2. It observed that, in all of genotypes there was a decrease in germination percentage due to salinity stress increment and maximum germination percentage was delayed. While in this experiment different genotypes had different response to the salinity stress. Among the safflower genotypes, Kose had the highest germination percentage and germination rate of 97.33% and 76.80 respectively. However, maximum reduction in germination percentage was observed at the highest level i.e., -1.5 MPa of NaCl (data not shown). At the highest salt level (-1.5 MPa), Kose produced maximum germination percentage and germination rate of all genotypes and they were considered as relatively tolerant.

Results of means comparison using Duncan multiple range test showed that germination percentage and germination rate decreased by increasing in osmotic potential and maximum germination rate and percentage were obtained at 0 Mpa level (control treatment). Some studies referred that stress can contribute to improve germination rate and seedling emergence in different plant species by increasing the expression of aquaporins (Gao *et al.*, 1999), enhancement of ATPase activity, RNA and acid phosphatase synthesis (Fu *et al.*, 1988), also by increase of amylases, proteases or lipases activity (Ashraf and Foolad, 2005).

Imposition of varying levels of NaCl significantly reduced all measured traits of all 6 investigated genotypes. Root length is one of the most important characters for salinity stress because roots are in contact with soil and absorb water from soil. For this reason, root length provides an important clue to the response of plants to salinity stress. A marked reduction in root length, shoot length and seedling length of all genotypes of safflower was observed due to salt stress.

Among the genotypes, the longest root length was commonly determined in genotypes FERAMAN, KM12, Kose and KM19 while SIRIN gave the shortest root length. Generally, increasing salinity levels decreased root length, and FERAMAN genotype exhibited the greater performance in respect of root length. Result of this study showed that, shoot length diminished with increasing salinity levels in all genotypes (Table 2). The highest and the lowest seedling length were observed in FERAMAN and SIRIN genotypes, respectively (Table 2).

Table 2. Mean comparison of main effects using Duncan multiple range test (at 5% probability level)

	Germination Percentage	Germination Rate	Root Length (mm)	Shoot Length (mm)	Seedling Length (cm)
Genotype					
KM12	77.44c	50.97c	4.02ab	5.08a	8.10a
KM19	70.33d	51.78bc	3.39ab	4.30b	6.69b
SIRIN	65.44d	44.13d	0.72c	0.31c	0.94c
FERAMAN	87.66b	53.55bc	4.18a	5.07a	8.25a
SINA	70.77cd	55.13b	3.21b	3.86b	6.08b
Kose	98.33a	77.80a	3.83ab	3.88b	6.71b
Salinity stress (MPa)					
0	82.85a	61.77a	4.25a	5.40a	8.65a
-0.3	79.51a	61.45ab	4.75a	5.71a	9.47a
-0.5	82.11a	58.26b	3.56b	4.09b	6.66b
-1	79.51a	52.70c	2.79c	3.01c	4.80c
-1.5	67.66b	43.63d	0.51d	0.38d	0.90d

Table 2. continued-

	Seed Vigour	Root/Shoot Length (mm)	Germination Index	Mean Germination Time (day)
Genotype				
KM12	580.48a	0.74ab	58.26c	3.03b
KM19	419.13b	0.78ab	52.86d	3.03b
SIRIN	63.17c	0.22c	46.60e	3.36a
FERAMAN	663.62a	0.73b	67.06b	2.937c
SINA	369.25b	0.76ab	54.60cd	2.88c
Kose	569.74a	1.13a	83.53a	2.36d
Salinity stress (MPa)				
0	652.90a	0.72a	65.88a	2.75c
-0.3	714.82a	0.68a	63.83ab	2.72c
-0.5	478.55b	0.72a	64.94ab	2.82c
-1	314.01c	0.75a	60.38b	2.98b
-1.5	60.87d	0.67a	47.38c	3.32a

Values in a column bearing different superscript are significantly different at 0.05 levels

The most effective levels in reducing these attributes were -1 and -1.5 MPa of NaCl (table 3). Best level of NaCl concentration in root length, shoot length, seedling length and seed vigour was -0.3 Mpa.

A significant inter-genotype variation was observed under salt stress. Of all genotypes, FERAMAN, KM12 and Kose produced highest seed vigour at all salt regimes, but lowest seed vigour was recorded in SIRIN while the remaining genotypes were moderate in this attribute.

Variation in the set of genotypes about root to shoot length ratio was not possible to discern at lower external salt levels, however, genotypes differed significantly at the two higher salt levels i.e. -1 and -1.5 MPa of NaCl (Table 3).

Table 3. Supplementary analysis of interaction effects.

Salinity Level (MPa)	Germination Percentage	Germination Rate	Root Length (mm)	Shoot Length (mm)	Seedling Length (cm)
0	624.57**	924.36**	4.08**	8.66**	20.84**
-0.3	981.69**	482.97**	11.01**	19.70**	59.31**
-0.5	365.93**	580.37**	6.21**	8.55**	28.48**
-1	626.21**	254.25**	3.08*	4.02**	13.23**
-1.5	396.99**	94.50**	0.19ns	0.44ns	1.03ns

Table 3. continued-

Salinity Level (MPa)	Seed Vigour	Root/Shoot Length (mm)	Germination Index	Mean Germination Time (cm)
0	25503**	0.15ns	749.28**	0.59**
-0.3	48837**	0.35ns	920.63**	0.48**
-0.5	23727**	0.16ns	472.98**	0.49**
-1	11225**	0.54**	551.92**	0.34**
-1.5	4044.98ns	0.51**	390.45**	0.36**

*, **, ns: significant at 5%, 1% level and not significant, respectively

In addition, it was clearly determined that there were no statistical differences between measured genotypes at high salinity levels (-1.5 MPa) for root length, shoot length, seedling length and seed vigour traits (Table 3).

Cluster analysis was done using the data for all measured traits at the highest salt level (-1.5 MPa), because this salt level was found very effective in discriminating the genotypes. Results of cluster analysis (Ward's minimum variance method) showed that genotypes Kose was found to be tolerant, while FERAMAN, SINA and SIRIN sensitive to salt (Fig. 1).




Figure1. Cluster analysis of safflower genotypes under -1.5 level of salinity stress using Ward's minimum variance method

Ajmal Khan and Weber (2006) found that, resistance to stress at germination stage and primary growth of seedling is independent from next growth stages and evaluation of stress tolerance need more experiment at next growth stages.

DISCUSSION

Screening of available germplasm of a crop is a feasible means of identifying salt tolerant genotypes or genotypes which could maintain a comparatively reasonable yield on salt affected soils (Ashraf & McNeilly, 1987). For the latter crops, it is advisable to assess degree of salt tolerance at each growth stage. In the present study, genotype Kose was found to be tolerant, while FERAMAN, SINA and SIRIN sensitive to salt. Ranking of the genotypes was done using the data for all measured traits at the highest salt level (-1.5 MPa), because this salt level was found very effective in discriminating the genotypes. These results can be related to some earlier studies in which genotypes identified as salt tolerant at the earlier growth stages showed tolerance when tested at the later growth stages.

Although a considerable magnitude of variation for salt tolerance was observed in a set of 6 available genotypes of safflower while screening them at germination stages, but a further study needs to be carried out to assess whether the genotypes marked as salt tolerant at the initial growth stages, maintain their degree of salt tolerance when tested as adult.

CONCLUSION

In the present study, salt stress adversely affected the germination percentage, germination rate, shoot length, root length, seedling length, and root to shoot length ratio, seed vigour, and germination index and mean germination time of all 6 genotypes of safflower and a significant variation in salt tolerance was observed among all the safflower.

Many researchers have been reported similar results (Demir, and Aril, 2003; Mauromicale and Licandro, 2002). Obviously, acceptable growth of plants in arid and semiarid lands which are under exposure of salinity stress is related to ability of seeds for best germination under unfavourable conditions, so necessity of evaluation of salinity resistance genotypes is important at primary growth stage. To find the best tolerant genotype to such conditions, taking all traits into account in this study, we found that Kose is the most resistant and FERAMAN, SINA and SIRIN are the most sensitive genotypes.

REFERENCES

- Ashraf M, Ahmad S. 1999. Exploitation of intra-specific genetic variation for improvement of salt (NaCl) tolerance in upland cotton (*Gossypium hirsutum* L.). *Hereditas*, 131: 253-256.
- Ashraf M, Fatima F. 1995. Responses of salt-tolerant and salt sensitive lines of safflower (*Carthamus tinctorius* L.) to salt stress. *Acta Physiol. Plant.*, 17(1): 61-70.
- Ashraf M, Foolad MR. 2005. Pre - sowing seed treatment - a shotgun approach to improve germination growth and crop yield under saline and none-saline conditions, *Advan. Agron.*, 88, 223-271.
- Ashraf M, McNeilly T. 1988. Variability in salt tolerance of nine spring wheat cultivars. *J. Agron. Crop Sci.*, 160: 14-21.
- Ashraf M, Waheed A. 1993. Screening of local/exotic accessions of lentil (*Lens culinaris* Medic.) for salt tolerance at two growth stages. *Plant Soil*, 128: 167-176.
- Ashraf M, Wahid S. 2000. Time-course changes in organic metabolites and mineral nutrients in germinating maize seeds under salt (NaCl) stress. *Seed Sci Technol* 28: 641-656
- Atak M, Kaya MD, Kaya G, Çıkılı Y, Çiftçi CY. 2006. Effects of NaCl on the germination, seedling growth and water uptake of triticale. *Turk. J. Agric. For.* 30: 39-47.
- Belkhdja R, Morales F, Abadia A, Gomez-Aparisi J, Abadia J. 1994. Chlorophyll fluorescence as a possible tool for salinity tolerance screening in barley (*Hordeum vulgare* L.). *Plant Physiol.*, 104: 667-673.
- Bernstein L. 1974. Crop growth and salinity. p 39-54. In: J. van Scdiffgard *Drainage for agriculture*. Agroomy. Monography, 17. ASA, Nadison, WI.
- Debez A, Ben Hamed K, Grignon C, Abdelly C. 2004. Salinity effects on germination, growth, and seed production of the halophyte *Cakile maritima*. *Plant Soil* 262: 179-189
- Fowler JL. 1991. Interaction of salinity and temperature on the germination of *Crambe*. *Agron. J.*, 83, 169-172.
- Francois LE, Bernstein L. 1964. Salt tolerance of safflower. *Agron. J.*, 56: 38-40.
- Ghorashy SR, Sionit N, Kheradnam M. 1972. Salt tolerance of safflower varieties (*Carthamus tinctorius* L.) during germination. *Agron. J.*, 64: 256-257.
- Golbashy M, Khavari Khorasani S, Ebrahimi M, Choukan R. 2010. Study of response of corn hybrids to limited irrigation. 11th Iranian Crop Science Congress Tehran, 24-26 July 2010. University of Shahid Beheshti. P. 218 (In Persian).
- Greenway H, Munns R. 1980. Mechanisms of salt tolerance in non-halophytes, *Annu. Rev. Plant Physiol.* 31: 149-190.
- Hair FF, Anderson RE, Tatham RL, Black WC. 1998. *Multivariate data analysis*, 5th Ed. Prentice-Hall Inc, Upple Saddle River, NJ, USA pp. 239-325.
- Jamil M, Lee DB, Jung KY, Ashraf M, Lee SC, Rha ES. 2006. Effect of salt (NaCl) stress on germination and early seedling growth of four vegetables species. *J. Cent. Eur. Agric.*, 7(2): 273-282.
- Janmohammadi M, Moradi Dezfuli P, Sharifzadeh F. 2008. Seed invigouration techniques to improve germination and early growth of inbred line of maize under salinity and drought stress. *Gen. Appl. Plant Physiology*.34 (3-4) .215-226.

- Kaya MD, Ipek A, Ozturk A. 2003. Effects of different soil salinity levels on germination and seedling growth of safflower (*Carthamus tinctorius* L.). *Turk. J. Agric.*, 27: 221-227.
- Kaya MD, Okçu G, Atak M, Çikili Y, Kolsarıcı O. 2006. Seed treatments to overcome salt and drought stress during germination in sunflower (*Helianthus annuus* L.). *Eur. J. Agron.* 24: 291-295.
- Kaya MD. 2009. The role of hull in germination and salinity tolerance in some sunflower (*Helianthus annuus* L.) cultivars. *African Journal of Biotechnology* . 8 (4), p: 597-600.
- Khaje-hosseini M, Powell AA, Bingham IJ. 2003. The interaction between salinity stress and seed vigour during germination of soybean seeds, *Seed Sci. Technol.*, 31, 715–725.
- Mohammed EM, Benbella M, Talouizete A. 2002. Effect of sodium chloride on sunflower (*Helianthus annuus* L.) seed germination. *Helia*, 25: 51-58.
- Murillo-Amador B, Lopez-Aguilar R, Kaya C, Larrinaga-Mayoral J, Flores Hernandez A. 2002. Comparative effect of NaCl and PEG on germination emergence and seedling growth of cowpea. *J. Agron. Crop Sci.*, 188, 235-247.
- Puppala N, Fowler JL, Poindexter L, Bhardavay HL. 1999. Evaluation of salinity tolerance of canola germination . p, 251-253. In. J. Jennick *Perpectives on new crops and new used*. ASHS Press, Alexandria,VA.
- Redmann RE, MQ QI, Belyk M. 1994. Growth of transgenic and standard Canola (*Brassica napus*) varieties in response to soil salinity. *Canadian Journal of Plant Science*, 74, 797-799.
- SAS. 2002. SAS/STAT® user's guide, SAS Institute, Cary, NC, USA. 9.1.
- Shannon MC. 1998. Adaptation of plant to salinity, *Adv. Agron.* 60: 75 119.
- Siddiqi E, Ashraf M, Aisha Akram N. 2007. Variation in seed germination and seedling growth in some diverse line of safflower (*Carthamus tinctorius* L.) under salt stress. *Pak. J. Bot.*, 39(6): 1937-1944.
- Ulfat M, Athar HR, Ashraf M, Akram NA, Jamil A. 2007. Appraisal of physiological and biochemical selection criteria for evaluation of salt tolerance in canola (*Brassica napus* L.). *Pak. J. Bot.*, 39(5): 1593-1608.
- Weiss EA. 1983. Oilseed crops. Longman Inc., New York.